

IN THE CASE OF A SEVERE NUCLEAR ACCIDENT RADIOACTIVE IODINE MAY BE RELEASED INTO THE ENVIRONMENT

Radioactive iodine do harm your health, especially children are at risk.

Iodine tablets (potassium iodine) prevent radioactive iodine from being absorbed and stored in your thyroid.

For further information:
www.jodblockade.de/en

This leaflet is based on the recommendation of the German Commission on Radiological Protection „Verwendung von Jodtabletten zur Jodblockade der Schilddrüse bei einem Notfall mit Freisetzung von radioaktivem Jod“ of 26. April 2018 (Jodmerkblätter)

Issuing point	Districts included
Fire station Bachl	Bachl, Birka, Scheuern, Stocka
Fire station Helchenbach	Aich, Asbach, Berg, Grub, Helchenbach, Höfel, Schöfthal, Kleinthalhof, Obermondsberg, Thalhof (bei Helchenbach), Weiherhof
Fire station Laaberberg	Alzhausen, Au, Kalteneck, Kronholzen, Laaber, Laaberberg, Mixmühle, Obergrünbach, Oberndorf, Schmiddorf, Untergrünbach
Fire station Oberbuch	Gressau, Oberbuch
Fire station Obereulenbach	Hirtdorf, Högetsing, Kiefermühle, Mordberg, Obereulenbach, See, Thalhof (bei Obereulenbach), Waselsdorf
Fire station Rohr i.NB	Rohr i.NB, Graben, Oberrohr, Reichenroith, Straßenhaus, Wallersdorf
Fire station Sallingberg	Sallingberg, Stegen, Ursbach

IMPRINT

Editor: Federal Ministry for the Environment, Nature Conservation and Nuclear Safety (BMU)
Division S II 5, P.O.Box 12 06 29, D-53048 Bonn
e-mail: SI15@bmu.bund.de | Internet: www.bmu.de

Editing: BMU, Division S II 5

Layout: :response (Gudrun Barth)

Photos: Jan Willem Geertsma (sxc.hu), :response

Date: June 2018



Federal Ministry
for the Environment, Nature Conservation
and Nuclear Safety



Intake of Iodine tablets

as protective measure in the event of a
severe accident in a nuclear power
plant

– information leaflet –

MAKE SURE THAT YOU TAKE THE CORRECT DOSAGE

The dosage is dependent on a person's age:



up to 1 month



from 1 month to 3 years



from 3 to 12 years



from 12 to 45 years and pregnant women




45 years and older

The risk of metabolism dysfunctions in thyroids increases with age. This so-called functional autonomy increases the risk of the iodine tablets side effects. Additionally, as people get older the risk of developing thyroid cancer caused by ionising radiation decreases considerably.

SUGGESTIONS FOR INTAKE

Either take the required amount by swallowing or by drinking it after dissolving the tablet in a liquid:



Dissolve the tablet in water



To facilitate the administration - especially for babies and children - dissolve the required amount in a drink, e.g. juice or tea.

Iodine tablets are a medicinal product. Please read the package insert carefully.

WHAT TO DO IN AN EMERGENCY?

- Pay close attention to the authorities announcements on radio and on television or to loud speaker announcements..
- These iodine tablets may only be taken on the explicit direction of the emergency response authority. Generally a single administration of the recommended dosage is sufficient. In exceptional cases the emergency response authorities may however recommend a further intake.
- The current timing of the tablet intake and the amount are very important.
- Prophylactic intake or other (higher) dosage do not reach a better protection - on the contrary - you could put your health at risk..
- Iodine tablets only provide protection against radioactive iodine. They do not provide protection against other radioactive substances.
- Therefore stay at home or seek shelter in windowless rooms.



Please take Iodine tablets only on the explicit direction of the emergency response